

Rhogogaster dryas (Benson). 1 ♀, 1 ♂, Crathie Wood, swept from *Populus tremula*, 23.6.84 (A.D.L.). The adults of this species, which is monophagous on aspen, are perhaps the most vividly coloured of all British sawflies. *R. dryas* is known from several English counties, but the only published Scottish records are of four males from Speyside (Benson, 1945; Woollatt, 1961).

Pseudodineura enslini (Hering). About a dozen leaf-mines were found on plants of *Trollius europaeus* growing in a meadow near Abergeldie Castle on 23.6.84. Although *Trollius* is locally common at other sites on the banks of the Dee, *P. enslini* could not be detected. The larvae were nearly full-grown when the mines were collected, and a few successfully spun cocoons. A single male emerged in August after having been "overwintered" in a refrigerator. In the British Isles, *enslini* has previously been recorded only from near Bettyhill in Sutherland, and near Grantown-on-Spey in Moray (Benson, 1958).

Pristiphora groenblomi (Lindqvist). 1 ♀, Morrone Birkwood NNR, Malaise Trap, 10.5.-1.6.84 (B. D. Batty). Previously in Britain only recorded from Aviemore (Benson, 1958). *P. groenblomi* is placed as a synonym of *coactula* (Ruthe) by Hellén (1975), but stronger evidence is required for such a synonymy because *groenblomi* is rather easily distinguished from the rest of the *coactula-breadalbanensis* complex.

References

- Benson, R. B., 1945. *Emphytus basalis* (Klug) and some other interesting British sawflies (Hym., Symphyta). *Ent.mon. Mag.* **81**: 101-103.
- Benson, R. B., 1958. Hymenoptera Symphyta. *Handbk. Ident. Br. Insects*, **5(2c)**: 139-252.
- Hellen, W., 1948. Mitteilungen uber einige Tenthredinoiden aus Ostfennoskandien. *Notulae Ent.* **28**: 40-46.
- Hellen, W., 1975. Die Nematinen Finnlands IV. *Notulae Ent.* **55**: 97-128.
- Woollatt, L. H., 1961. Sawflies (Hym., Symphyta) collected by Mr. J. E. Collin and Mr. E. C. M. d'Assis-Fonseca in the Scottish Highlands, May and June, 1959. *Ent. mon. Mag.* **97**: 87-88.

PERIDEA ANCEPS GOEZE: GREAT PROMINENT, IN LATE JULY. — On 28 July 1984 I collected a male *Peridea anceps* at M.V. light at the Centre for Alternative Technology, Montgomeryshire. This species usually appears in spring from April to June and so a late July record is not only unusual but raises the question of either a second brood (unlikely) or of an immigrant (equally unlikely). Most probably the moth emerged late or somehow managed to live a long adult life. — DENIS F. OWEN, 66 Scraptoft Lane, Leicester LE5 1HU.